



SHORT REPORT

Open Access

Elevated serum fibroblast growth factor 21 levels correlate with immune recovery but not mitochondrial dysfunction in HIV infection

Brendan Al Payne^{1,2*}, David Ashley Price² and Patrick F Chinnery¹

Abstract

Background: Anti-retroviral treated HIV-infected patients are at risk of mitochondrial toxicity, but non-invasive markers are lacking. Serum FGF-21 (fibroblast growth factor 21) levels correlate strongly with muscle biopsy findings in inherited mitochondrial disorders. We therefore aimed to determine whether serum FGF-21 levels correlate with muscle mitochondrial dysfunction in HIV-infected patients.

Findings: We performed a cross-sectional study of anti-retroviral treated HIV-infected subjects (aged 29 – 71 years, $n = 32$). Serum FGF-21 levels were determined by quantitative ELISA. Cellular mitochondrial dysfunction was assessed by COX (cytochrome *c* oxidase) histochemistry of lower limb skeletal muscle biopsy. Serum FGF-21 levels were elevated in 66% of subjects. Levels correlated significantly with current CD4 lymphocyte count ($p = 0.042$) and with total CD4 count gain since initiation of anti-retroviral therapy ($p = 0.016$), but not with the nature or duration of past or current anti-retroviral treatment. There was no correlation between serum FGF-21 levels and severity of the muscle mitochondrial (COX) defect.

Conclusions: Serum FGF-21 levels are a poor predictor of muscle mitochondrial dysfunction in contemporary anti-retroviral treated patients. Serum FGF-21 levels are nevertheless commonly elevated, in association with the degree of immune recovery, suggesting a non-mitochondrial metabolic disturbance with potential implications for future comorbidity.

Keywords: HIV, Anti-retroviral therapy, Highly active, Fibroblast growth factor 21, Mitochondria

Findings

Mitochondrial dysfunction is a well-described complication of anti-retroviral therapy [1-7]. It is most strongly associated with several of the older nucleoside analogue reverse transcriptase inhibitors (NRTIs): zidovudine (AZT), stavudine (d4T), didanosine (ddI), and zalcitabine (ddC). Although these drugs are no longer in common usage in industrialised countries, there are nevertheless large numbers of patients who have had extensive prior exposure to these drugs, and some remain in common usage in developing countries. We have recently demonstrated that patients with previous exposure to these NRTIs may have

persistent cellular mitochondrial COX (cytochrome *c* oxidase) defects in skeletal muscle, consequent on an NRTI-induced accumulation of somatic (acquired) mitochondrial DNA (mtDNA) mutations [8].

Non-invasive measures of mitochondrial damage would be very valuable in the HIV clinic, both for the diagnosis of anti-retroviral associated mitochondrial dysfunction and the serial monitoring of such patients. The determination of mtDNA content in peripheral blood mononuclear cells (PBMCs) has previously been proposed as such a measure [5,9-11]. This consideration arises from the fact that the mitochondrially-toxic NRTIs (as listed above) cause a reduction in cellular mtDNA content (depletion) during therapy [2,3,6,10,12-16]. However, modern N(t) RTIs such as tenofovir (TDF) and abacavir (ABC) do not cause mtDNA depletion [17], and as a result, mtDNA levels return to normal with a switch away

* Correspondence: brendan.payne@ncl.ac.uk

¹Institute of Genetic Medicine, Newcastle University, Central Parkway, Newcastle-upon-Tyne NE1 3BZ, UK

²Department of Infection and Tropical Medicine, Royal Victoria Infirmary, Newcastle-upon-Tyne NE1 4LP, UK

from a mitochondrially-toxic NRTI. Thus, measuring mtDNA levels is not a useful measure of on-going mitochondrial dysfunction due to an NRTI exposure in the distant past.

In contrast, FGF-21 (fibroblast growth factor 21) has recently been proposed as a valuable serum measure in inherited mitochondrial disease [18]. In these patients, serum FGF-21 levels showed a very strong correlation with mitochondrial dysfunction on skeletal muscle biopsy, as determined by the percentage of cells expressing a COX defect [18]. FGF-21 is thought to regulate mitochondrial activity and enhance oxidative capacity, mediated via PGC-1 α (peroxisome proliferator-activated receptor gamma co-activator 1-alpha) expression [19]. To date, one study has assessed serum FGF-21 in HIV infection, and demonstrated elevated levels [20]. Given the recently described association between serum FGF-21 elevation and muscle COX defects in inherited mitochondrial disorders [18], and the recent observation of significant COX defects in long-term anti-retroviral treated HIV-infected patients [8], we speculated that muscle mitochondrial dysfunction might also drive the FGF-21 elevation in anti-retroviral treated HIV infection.

Patient characteristics

All subjects gave informed written consent for participation, and the study was approved by local research ethics committee (Newcastle and North Tyneside Research Ethics Committee). We performed a cross-sectional study of adult HIV-1 infected patients, receiving ambulatory care at one of two specialist clinics in Newcastle-upon-Tyne, UK. Patients with current active hepatitis B or C co-infection, known inherited or non-HIV-associated neuromuscular disease, and diabetes mellitus were excluded. No subjects were clinically obese (BMI >30). 32 HIV-infected subjects participated, of whom 81% were male. 84% were of white Caucasian ethnicity and the remainder black African. Mean age was 48.7 years, with age range of 29–71 years. Mean duration of diagnosed HIV infection was 10.8 years. Mean current CD4 lymphocyte count was 663 cells/ μ l, and 61% of subjects had nadir CD4 count of <200 cells/ μ l. All subjects were currently receiving combination anti-retroviral therapy, with a mean duration of treatment of 9.2 years. 97% of patients had fully suppressed HIV plasma viral load (<50 HIV-1 RNA copies/ml). 81% of subjects were receiving a non-nucleoside reverse transcriptase inhibitor (NNRTI) and 22% a ritonavir-boosted protease inhibitor (PI). Regarding past (lifetime) NRTI treatment experience, 72% of patients had a history of AZT exposure, and 25% had a history of prior d-drug (dideoxynucleoside analogue) exposure. Characteristics of individual subjects are shown in Table 1.

FGF-21 determination

Serum FGF-21 levels were determined by quantitative ELISA (BioVendor, Brno, Czech Republic), performed in triplicate, and normalised by log₁₀ transformation. A serum FGF-21 level of <200 pg/ml was considered as normal in keeping with recent data [18]. Statistical analyses were performed in SPSS 19, using student's t-test to compare binary variables and Pearson's correlation coefficient (*r*) to examine the relationship between log₁₀ serum FGF-21 levels and continuous variables. Twenty-one of 32 subjects (66%) had serum FGF-21 levels greater than the normal range, with four being very elevated (>800 pg/ml). On univariate analysis, serum FGF-21 levels were positively correlated with current CD4 lymphocyte count (*r* = 0.36, *p* = 0.042), but more strongly correlated with total CD4 cell count gain since initiation of anti-retroviral therapy (current minus nadir) (*r* = 0.45, *p* = 0.016) (Figure 1). In addition, plasma glucose levels correlated with serum FGF-21 levels, although this did not quite reach statistical significance (*r* = 0.34, *p* = 0.06, Figure 2), whereas as serum lipids and liver function did not. No other demographic or treatment variables were significantly associated with serum FGF-21 levels, including the nature of current or prior anti-retroviral therapy (Table 2). FGF-21 levels did not differ significantly between patients with or without clinical lipodystrophy syndrome. Only CD4 lymphocyte count gain was independently associated with serum FGF-21 levels on multivariate linear regression analysis (*p* = 0.016).

Skeletal muscle mitochondrial histochemistry

COX histochemistry was performed on cryo-sections obtained from lower limb skeletal muscle biopsies on 31 of the 32 subjects (biopsy data for one subject was not analysable). Results of 22 of these biopsies have been reported in our previous work [8], whereas the remaining 9 have not. COX contains respiratory chain subunits encoded by the mitochondrial genome, and fibres stain brown (positive) in the presence of intact respiratory chain activity (Figure 3). Proportional COX defect was determined by counting \geq 500 fibres per biopsy, and normalised by log₁₀ transformation. There was no correlation between serum FGF-21 levels and percentage COX defects on biopsy (*r* = -0.02, *p* = 0.9, Figure 4).

Discussion

We have shown that serum FGF-21 levels are frequently elevated in contemporary anti-retroviral treated HIV-infected patients, but do not correlate with the severity of muscle mitochondrial (COX) defect. In contrast, a previous study has shown a very strong correlation between these parameters in patients with inherited mitochondrial disorders [18]. Ours is the first study to

Table 1 Patient characteristics

Age (y)	Gender	Ethnicity	Duration of diagnosed HIV (mo)	ART duration (mo)	Nadir CD4 count (cells/uL)	Current CD4 count (cells/uL)	LDS	ART (current)	ART (lifetime)	COX defect (%)	Serum FGF-21 (pg/mL)
71	M	WB	130	130	UK	530	Y	TDF FTC EFV	ddi AZT 3TC EFV TDF FTC	3.0%	232
48	F	BA	100	99	10	487	N	TDF FTC NVP	AZT 3TC EFV TDF FTC NVP	0.1%	>1920
34	F	WB	88	86	218	1121	Y	ABC 3TC NVP	AZT 3TC NVP ABC	0.0%	560
55	F	BA	64	27	112	426	N	TDF FTC AZT DRV/r	TDF FTC LPV/r AZT DRV/r	0.8%	164
43	M	BA	87	87	152	306	N	TDF FTC EFV	AZT 3TC EFV TDF FTC	0.8%	20
42	M	WB	185	147	150	636	N	AZT 3TC EFV	AZT 3TC NVP	0.0%	342
63	M	WB	97	97	169	870	N	ABC 3TC EFV	AZT 3TC EFV ABC	0.0%	204
29	M	WB	84	32	197	401	N	TDF FTC EFV	TDF FTC EFV	0.0%	809
63	M	WB	238	221	NA	438	N	ABC 3TC NVP	AZT ddi d4T 3TC ddC IDV NVP ABC	2.2%	440
62	M	WB	63	62	56	190	N	TDF FTC NVP	TOP FTC NVP	0.2%	156
52	M	WB	225	225	120	728	Y	TDF FTC ATV/r	AZT ddC ddi 3TC d4T SQV NVP IDV NFV ABC TDF LPV/r FTC ATV/r	1.3%	328
36	M	WB	139	138	197	627	N	AZT 3TC EFV	AZT 3TC EFV TDF FTC	0.3%	214
51	M	WB	190	183	10	747	N	ABC TDF NVP	AZT ddi d4T 3TC RTV NVP IDV ddC ABC ATV/r TDF	4.9%	252
33	F	WB	96	95	83	1289	N	TDF FTC EFV	AZT 3TC EFV TDF FTC	0.0%	177
48	M	WB	102	100	259	1329	Y	AZT 3TC EFV	AZT 3TC EFV	0.4%	409
51	M	WB	145	144	151	421	N	AZT 3TC NVP	AZT 3TC NVP	1.4%	254
66	M	WB	71	26	287	455	N	TDF FTC EFV	TDF FTC EFV	11.2%	470
46	M	WB	158	157	250	1452	N	TDF FTC EFV	AZT 3TC IDV EFV ABC TDF FTC	1.4%	1218
61	M	WB	116	113	NA	498	Y	TDF FTC NVP	AZT 3TC EFV NVP TDF FTC	2.4%	182-
30	M	WB	88	23	283	661	N	TDF FTC DRV/r	TDF FTC EFV DRV/r	0.1%	207
62	M	WB	284	202	NA	422	N	ABC NVP LPV/r	SQV AZT ddC 3TC d4T ddi IDV ABC NVP NFV LPV/r	0.8%	92
45		WB	159	158	176	660	N	TDF FTC NVP	AZT 3TC IDV NVP TDF FTC	0.0%	60
54	M	WB	79	38	244	638	N	TDF FTC DRV/r	TDF FTC EFV DRV/r	3.4%	178
52	M	WB	166	164	0	662	Y	TDF FTC NVP	AZT d4T IDV NFV SQV 3TC NVP ddi TDF FTC	2.8%	525
51	M	WB	243	171	327	539	Y	TDF FTC EFV	AZT ddi RTV NFV TDF FTC EFV	1.5%	231
35	F	BA	62	25	380	638	N	TDF FTC EFV	TOP FTC EFV	0.0%	82
53	M	WB	NA	48	301	804	N	TDF FTC EFV	TDF FTC EFV	NA	490
36	M	WB	130	130	18	898	N	TDF FTC ATV/r	AZT 3TC EFV TDF FTC ATV/r	0.0%	530
48	M	WB	53	14	332	443	N	TDF FTC EFV	TDF FTC EFV	0.0%	56

Table 1 Patient characteristics (Continued)

52	F	BA	83	81	17	485	Y	TDF FTC EFV	AZT 3TC EFV TDF FTC	0.7%	224
38	M	WB	129	128	4	761	Y	TDF FTC EFV	AZT 3TC EFV TDF FTC	0.0%	935
47	M	WB	183	164	305	668	Y	ABC RAL ATV/r	d4T 3TC NVP ddi IDV ABC ATV/r RAL	9.8%	114

Summary characteristics of individual HIV-infected subjects. (WB, white British; BA, black African; ART, anti-retroviral therapy; LDS, lipodystrophy syndrome; AZT, zidovudine; d4T, stavudine; ddi, didanosine; ddC, zalcitabine; 3TC, lamivudine; ABC, abacavir; TDF, tenofovir; FTC, emtricitabine; EFV, efavirenz; NVP, nevirapine; ATV, atazanavir; DRV, darunavir; LPV, lopinavir; SQV, saquinavir; NFV, nelfinavir; IDV, indinavir; RTV, ritonavir at therapeutic dose; /r, ritonavir at pharmacokinetic boosting dose; RAL, raltegravir; COX, cytochrome c oxidase; FGF-21, fibroblast growth factor 21; NA, not available).




Figure 1 Correlation of serum FGF-21 levels with immune reconstitution. Correlation of \log_{10} serum FGF-21 (fibroblast growth factor 21) levels in HIV-infected subjects and CD4 lymphocyte count gain on treatment (current minus nadir) ($r = 0.45$, $p = 0.016$). (* Serum FGF-21 >1920 pg/ml, the upper limit of quantitation of the assay).

attempt to link serum FGF-21 levels with biopsy-proven mitochondrial defects in HIV-infected patients. What is the reason for this apparent discrepancy in findings? Firstly, the prior study demonstrating serum FGF-21 elevation in mitochondrial disease included a large number of patients with childhood-onset disease. Such patients typically have very severe muscle COX defects (affecting up to ~60% of fibres). In contrast,




Figure 2 Correlation of serum FGF-21 levels with plasma glucose. Correlation of \log_{10} serum FGF-21 (fibroblast growth factor 21) levels in HIV-infected subjects and random plasma glucose concentration ($r = 0.34$, $p = 0.06$). (* Serum FGF-21 >1920 pg/ml, the upper limit of quantitation of the assay).

Table 2 Associations of serum FGF-21 levels

(a)			
Variable	(n)	\log_{10} Serum FGF-21, mean (SD)	p value
Gender	Male (26)	2.39 (0.40)	0.71
	Female (6)	2.46 (0.48)	
Ethnicity	Caucasian (27)	2.44 (0.33)	0.27
	Black African (5)	2.21 (0.72)	
Current ART	PI (7)	2.29 (0.26)	0.43
	No PI (25)	2.43 (0.44)	
Lifetime ART	d-drugs (8)	2.38 (0.26)	0.86
	No d-drugs (24)	2.41 (0.45)	
	AZT (23)	2.44 (0.42)	
Lipodystrophy	No AZT (9)	2.30 (0.39)	0.41
	Yes (10)	2.50 (0.27)	
Lipid-lowering therapy	No (22)	2.36 (0.46)	0.39
	Yes (8)	2.32 (0.52)	
	No (24)	2.43 (0.38)	0.53

(b)			
Variable		Correlation coefficient (r)	p value
Age		-0.07	0.69
Duration of diagnosed HIV infection		-0.08	0.67
Duration of lifetime ART	Total	0.12	0.50
	d-drug	0.01	0.97
	AZT	0.13	0.47
CD4 lymphocyte count	Nadir	-0.29	0.14
	Current	0.36	0.042
CD4 count gain	(Current minus nadir)	0.45	0.016
Serum ALT		0.27	0.14
Plasma glucose		0.34	0.06
Serum lipids	Total cholesterol	-0.03	0.87
	HDL cholesterol	0.02	0.90
	Non-HDL cholesterol	-0.06	0.75
Mitochondrial histochemistry	COX defect (\log_{10})	-0.02	0.91

(a) binary variables; (b) continuous variables. (PI, protease inhibitor; AZT, zidovudine; d-drug, dideoxynucleoside analogue; ART, anti-retroviral therapy; ALT, alanine transaminase; HDL, high density lipoprotein). Bold type indicates statistically significant associations.




Figure 3 COX histochemistry. Example of mitochondrial COX/SDH (cytochrome c oxidase/succinate dehydrogenase) histochemistry on lower limb skeletal muscle biopsy of an anti-retroviral treated HIV-infected patient. Normal (COX positive) fibres stain brown, whereas COX deficient fibres counterstain blue due to preserved SDH activity.

patients with late-onset inherited mitochondrial disorders typically have more modest COX defects, comparable with those seen in our HIV-infected patients (up to ~10% of fibres). The fact that some patients in our study with a biopsy COX defect of >5% of fibres had relatively normal FGF-21 levels suggests that this serum measure is not particularly sensitive for mild to moderate muscle mitochondrial defects. Secondly, the markedly abnormal serum FGF-21 levels seen in some patients with no significant COX defect suggest a non-mitochondrial origin, as has been observed in other




Figure 4 Correlation of serum FGF-21 levels and mitochondrial defects. Correlation of \log_{10} serum FGF-21 (fibroblast growth factor 21) levels in HIV-infected subjects and percentage COX (cytochrome c oxidase) defect on lower limb skeletal muscle biopsy ($r = -0.02$, $p = 0.9$). (* Serum FGF-21 >1920 pg/ml, the upper limit of quantitation of the assay).

metabolic disorders [21-23]. In the only previous study of FGF-21 levels in HIV infection, the authors found associations of FGF-21 levels with obesity, glycaemia, dyslipidaemia and liver dysfunction, in line with literature from HIV-uninfected patients [24]. In our study, we specifically excluded diabetic and obese subjects (as we wished to maximise the likelihood of detecting any association with NRTI-induced mitochondrial dysfunction). Interestingly however, the strongest predictor of serum FGF-21 levels seen in our study was a novel association with total CD4 lymphocyte count gain. This is an intriguing finding. It is plausible that patients who have low nadir CD4 lymphocyte counts may experience more profound metabolic changes as they undergo immune reconstitution on anti-retroviral therapy, switching from a catabolic state to an excessively anabolic state associated with a 'return to health'. This association with CD4 count gain should be further explored by longitudinal study.

In conclusion, serum FGF-21 levels do not appear to be a sensitive or specific marker of muscle mitochondrial dysfunction in contemporary anti-retroviral treated patients. Nevertheless they are commonly elevated in association with immune recovery. As serum FGF-21 levels in the HIV-uninfected population are elevated in conditions associated with increased cardiovascular risk, it is very plausible that serum FGF-21 elevation in anti-retroviral treated HIV infection may also be a marker of an adverse metabolic risk in this patient group. Given the known increase in cardiovascular disease in anti-retroviral treated patients [25], the prognostic significance of our findings merits further research.

Competing interests

All authors confirm that they have no relevant competing interests.

Authors' contributions

BP conceived the study, performed the assays and drafted the manuscript. DAP conceived and helped coordinate the study. PFC conceived the study and drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to acknowledge Alison Hague for assistance with the FGF-21 ELISA.

Funding

Medical Research Council (UK) (BP); Newcastle National Institute for Health Research (NIHR) Biomedical Research Centre in Ageing (BP, PFC); Medical Research Council (UK) Centre for Translational Muscle Disease (PFC); Wellcome Trust (084980/Z/08/Z & 096919Z/11/Z, PFC).

Received: 12 July 2013 Accepted: 14 November 2013

Published: 19 November 2013

References

- Dalakas MC, Illa I, Pezeshkpour GH, Laukaitis JP, Cohen B, Griffin JL: **Mitochondrial myopathy caused by long-term zidovudine therapy.** *N Engl J Med* 1990, **322**:1098–1105.
- Arnaudo E, Dalakas M, Shanske S, Moraes CT, DiMauro S, Schon EA: **Depletion of muscle mitochondrial DNA in AIDS patients with zidovudine-induced myopathy.** *Lancet* 1991, **337**:508–510.
- Shikuma CM, Hu N, Milne C, Yost F, Waslien C, Shimizu S, Shiramizu B: **Mitochondrial DNA decrease in subcutaneous adipose tissue of HIV-infected individuals with peripheral lipodystrophy.** *Aids* 2001, **15**:1801–1809.
- Zaera MG, Miro O, Pedrol E, Soler A, Picon M, Cardellach F, Casademont J, Nunes V: **Mitochondrial involvement in antiretroviral therapy-related lipodystrophy.** *Aids* 2001, **15**:1643–1651.
- van der Valk M, Casula M, Weverling GJ, van Kuijk K, van Eck-Smit B, Hulsebosch HJ, Nieuwkerk P, van Eeden A, Brinkman K, Lange J, et al: **Prevalence of lipodystrophy and mitochondrial DNA content of blood and subcutaneous fat in HIV-1-infected patients randomly allocated to zidovudine- or stavudine-based therapy.** *Antivir Ther* 2004, **9**:385–393.
- Walker UA, Bauerle J, Laguno M, Murillas J, Mauss S, Schmutz G, Setzer B, Miquel R, Gatell JM, Mallolas J: **Depletion of mitochondrial DNA in liver under antiretroviral therapy with didanosine, stavudine, or zalcitabine.** *Hepatology* 2004, **39**:311–317.
- Miro O, Lopez S, Pedrol E, Rodriguez-Santiago B, Martinez E, Soler A, Milinkovic A, Casademont J, Nunes V, Gatell JM, Cardellach F: **Mitochondrial DNA depletion and respiratory chain enzyme deficiencies are present in peripheral blood mononuclear cells of HIV-infected patients with HAART-related lipodystrophy.** *Antivir Ther* 2003, **8**:333–338.
- Payne BA, Wilson IJ, Hately CA, Horvath R, Santibanez-Koref M, Samuels DC, Price DA, Chinnery PF: **Mitochondrial aging is accelerated by anti-retroviral therapy through the clonal expansion of mtDNA mutations.** *Nat Genet* 2011, **43**:806–810.
- Chiappini F, Teicher E, Saffroy R, Pham P, Falissard B, Barrier A, Chevalier S, Debuire B, Vittecoq D, Lemoine A: **Prospective evaluation of blood concentration of mitochondrial DNA as a marker of toxicity in 157 consecutively recruited untreated or HAART-treated HIV-positive patients.** *Lab Invest* 2004, **84**:908–914.
- Cote HC, Brumme ZL, Craib KJ, Alexander CS, Wynhoven B, Ting L, Wong H, Harris M, Harrigan PR, O'Shaughnessy MV, Montaner JS: **Changes in mitochondrial DNA as a marker of nucleoside toxicity in HIV-infected patients.** *N Engl J Med* 2002, **346**:811–820.
- Montaner JS, Cote HC, Harris M, Hogg RS, Yip B, Chan JW, Harrigan PR, O'Shaughnessy MV: **Mitochondrial toxicity in the era of HAART: evaluating venous lactate and peripheral blood mitochondrial DNA in HIV-infected patients taking antiretroviral therapy.** *J Acquir Immune Defic Syndr* 2003, **34**(Suppl 1):S85–S90.
- Cherry CL, Gahan ME, McArthur JC, Lewin SR, Hoy JF, Wesselingh SL: **Exposure to dideoxynucleosides is reflected in lowered mitochondrial DNA in subcutaneous fat.** *J Acquir Immune Defic Syndr* 2002, **30**:271–277.
- Hoschele D: **Cell culture models for the investigation of NRTI-induced mitochondrial toxicity. Relevance for the prediction of clinical toxicity.** *Toxicol In Vitro* 2006, **20**:535–546.
- Dalakas MC, Semino-Mora C, Leon-Monzon M: **Mitochondrial alterations with mitochondrial DNA depletion in the nerves of AIDS patients with peripheral neuropathy induced by 2'3'-dideoxycytidine (ddC).** *Lab Invest* 2001, **81**:1537–1544.
- Hammond E, Nolan D, James I, Metcalf C, Mallal S: **Reduction of mitochondrial DNA content and respiratory chain activity occurs in adipocytes within 6–12 months of commencing nucleoside reverse transcriptase inhibitor therapy.** *Aids* 2004, **18**:815–817.
- Lim SE, Copeland WC: **Differential incorporation and removal of antiviral deoxynucleotides by human DNA polymerase gamma.** *J Biol Chem* 2001, **276**:23616–23623.
- Birkus G, Hitchcock MJ, Cihlar T: **Assessment of mitochondrial toxicity in human cells treated with tenofovir: comparison with other nucleoside reverse transcriptase inhibitors.** *Antimicrob Agents Chemother* 2002, **46**:716–723.
- Suomalainen A, Elo JM, Pietilainen KH, Hakonen AH, Sevastianova K, Korpela M, Isohanni P, Marjavaara SK, Tyni T, Kiuru-Enari S, et al: **FGF-21 as a biomarker for muscle-manifesting mitochondrial respiratory chain deficiencies: a diagnostic study.** *Lancet Neurol* 2011, **10**:806–818.
- Moyers JS, Shiyanova TL, Mehrbod F, Dunbar JD, Noblitt TW, Otto KA, Reifel-Miller A, Kharitononkov A: **Molecular determinants of FGF-21 activity-synergy and cross-talk with PPARgamma signaling.** *J Cell Physiol* 2007, **210**:1–6.
- Affandi JS, Price P, Imran D, Yuniastuti E, Djauzi S, Cherry CL: **Can we predict neuropathy risk before stavudine prescription in a resource-limited setting?** *AIDS Res Hum Retroviruses* 2008, **24**:1281–1284.
- Chen WW, Li L, Yang GY, Li K, Qi XY, Zhu W, Tang Y, Liu H, Boden G: **Circulating FGF-21 levels in normal subjects and in newly diagnose patients with Type 2 diabetes mellitus.** *Exp Clin Endocrinol Diabetes* 2008, **116**:65–68.
- Dushay J, Chui PC, Gopalakrishnan GS, Varela-Rey M, Crawley M, Fisher FM, Badman MK, Martinez-Chantar ML, Maratos-Flier E: **Increased fibroblast growth factor 21 in obesity and nonalcoholic fatty liver disease.** *Gastroenterology* 2010, **139**:456–463.
- Zhang X, Yeung DC, Karpisek M, Stejskal D, Zhou ZG, Liu F, Wong RL, Chow WS, Tso AW, Lam KS, Xu A: **Serum FGF21 levels are increased in obesity and are independently associated with the metabolic syndrome in humans.** *Diabetes* 2008, **57**:1246–1253.
- Domingo P, Gallego-Escuredo JM, Domingo JC, Gutierrez Mdel M, Mateo MG, Fernandez I, Vidal F, Giral M, Villarroya F: **Serum FGF21 levels are elevated in association with lipodystrophy, insulin resistance and biomarkers of liver injury in HIV-1-infected patients.** *AIDS* 2010, **24**:2629–2637.
- Friis-Moller N, Sabin CA, Weber R, D'Arminio Monforte A, El-Sadr WM, Reiss P, Thiebaut R, Morfeldt L, De Wit S, Pradier C, et al: **Combination antiretroviral therapy and the risk of myocardial infarction.** *N Engl J Med* 2003, **349**:1993–2003.

doi:10.1186/1742-6405-10-27

Cite this article as: Payne et al.: Elevated serum fibroblast growth factor 21 levels correlate with immune recovery but not mitochondrial dysfunction in HIV infection. *AIDS Research and Therapy* 2013 **10**:27.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

