BRIEF COMMUNICATION

Presence of tuberculosis symptoms among HIV-positive men who have sex with men (MSM) in Zimbabwe

Munyaradzi Mapingure¹, Innocent Chingombe¹, Tafadzwa Dzinamarira¹, Brian Moyo², Chesterfield Samba³, Delight Murigo³, Owen Mugurungi², Elliot Mbunge⁴, Rutendo Birri Makota⁵, Grant Murewanhema⁶ and Godfrey Musuka^{7*}

Abstract

We conducted secondary data analysis using a biobehavioral survey dataset of 1538 MSM from Zimbabwe. Survey participants were screened for the four symptoms suggestive of tuberculosis infection using the WHO TB screening algorithm. Results: All participants experienced at least one symptom suggestive of tuberculosis. 40% of HIV-positive MSM reported having had a cough in the last month and 13% of them experienced unexpected weight loss. The prevalence of experiencing any of the four TB symptoms amongst HIV-positive MSM was 23%.

Contribution

There is an urgent need for active TB case finding and treatment amongst HIV-positive MSM in Zimbabwe. Clinicians will need to ensure that MSM who need TB testing receive it timeously.

in this article, unless otherwise stated in a credit line to the data.

Keywords HIV, Tuberculosis symptoms, Case finding, MSM

gmusuka@3ieimpact.org ¹ICAP in Zimbabwe, Harare, Zimbabwe ²AIDS and TB Programmes, Ministry of Health and Child Care, Harare,

Zimbabwe

*Correspondence:

Godfrev Musuka

³GALZ, Harare, Zimbabwe

⁴Department of Computer Science, Faculty of Science and Engineering, University of Eswatini, Kwaluseni, Eswatini

⁵Department of Biological Sciences and Ecology, University of Zimbabwe, Harare, Zimbabwe

⁶Unit of Obstetrics and Gynaecology, Faculty of Medicine and Health Sciences, University of Zimbabwe, Harare, Zimbabwe

⁷International Initiative for Impact Evaluation, Harare, Zimbabwe

Background

© The Author(s) 2024. Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use,

sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available

Despite being both preventable and treatable, TB remains an important public health problem in all countries and age groups [1]. Globally and in sub-Saharan Africa, tuberculosis is a major cause of mortality and morbidity [2, 3]. Being HIV-positive without viral load suppression is a predisposing factor for TB infection, amongst MSM. TB prevalence surveys in resource-poor countries consistently show a high burden of patients with TB disease who remain undiagnosed and are thus infectious to others [4]. Improving international case detection for TB also contributes to achieving health-related global goals [5]. HIV prevalence is declining in Zimbabwe but remains one of the highest in sub-Saharan Africa at 11.6% among the







general population [6] and is higher at 23.4% among men who have sex with men (MSM) [7].

Eliciting of symptoms is an essential step in the diagnosis and care of presumptive MSM tuberculosis patients [8]. MSM are among the critical populations identified to be at higher risk of tuberculosis infection and disease [9]. However, there is limited TB data among MSM [10, 11].

Methods

Data from our study on HIV and STI biobehavioral surveys (BBS) among MSM in Zimbabwe [7] was used to determine the prevalence of tuberculosis (TB) symptoms amongst HIV-positive MSM. Data were collected from 1538 individual MSM in the Zimbabwe cities of Harare and Bulawayo.

The study was a cross-sectional BBS using respondent-driven sampling (RDS). RDS is a peer-referral sampling methodology designed for data collection among hard-to-reach populations. Consenting participants completed a questionnaire on sociodemographics, sexual identity and history, sexual behaviour, mental health, alcohol and drug use, HIV testing services, stigma and violence, social cohesion, health services and TB. As part of the study, participants were screened for the four symptoms suggestive of tuberculosis infection using the WHO TB screening algorithm, and to our knowledge, this is the first and only BBS in Zimbabwe to have done so. After being tested for HIV, participants were also screened for tuberculosis (TB) symptoms. Participants who screened positive for TB symptoms were offered the option of being driven that day to a designated MSM-friendly referral healthcare facility that offers MSM-friendly HIV and other related services. Additionally, a peer escort to accompany them was provided. If the participant declined, they were given a referral card for the clinic that offers MSM-friendly HIV and related services. STATA statistical package version 17 was used for statistical analysis. The study received ethical clearance from the national institutional review board, the Medical Research Council of Zimbabwe # (MRCZ/A/2156).

Results

In terms of demographics, the HIV-positive MSM were older than the negative, with Mean (Standard deviations) of 31.8 (9.2) and 26.4 (7.9) years respectively, p=0.001. The majority (>96%) were Black Africans, who had primarily attained (>70%) secondary education. Close to 40% were unemployed, see Table 1. All participants experienced at least one symptom suggestive of tuberculosis, with prevalences of 7%, 4%, 7%, and 1% for persistent cough, night sweats, unexpected weight loss and fever respectively. Having had a cough

in the last month was higher among HIV-positive MSMs compared to those not affected with HIV, 40% vs. 5%, p=0.001. The same was the case with experiencing unexpected weight loss, 13% vs. 6%, p=0.001 and experiencing a fever in the last month, 3% vs. 1%, p=0.037. The prevalence of experiencing any of the four TB symptoms was 13%, and this prevalence was higher among the HIV-positive MSM compared to those who are not infected with HIV, 23% vs. 11%, p=0.001. Three per cent reported that they had been exposed to a person with TB in the last 12 months.

Discussions

A systematic review by Divala and others [8] shows that invariably, tuberculosis symptom screening is a critical entry point for the diagnosis of the disease [12-15]. Additionally, there is a need for more accurate screening tools, that remain accessible and low-cost for countries with high tuberculosis burden.

Some individuals wait for the spontaneous resolution of TB symptoms, and most do not seek healthcare help [15]. The biological and epidemiological interconnectedness of TB and HIV has generated a new form of stigma, TB-HIV stigma, which sometimes makes people afraid to go for TB symptoms as this would be perceived to be HIV-related TB.

We note that there is limited information on TB among critical populations such as MSM, and the findings in this letter are crucial for understanding TB symptoms among MSM. There is a need for ongoing campaigns to encourage people to seek healthcare help when experiencing TB symptoms so that we do not have infectious patients who remain undiagnosed and are thus contagious to others. Targeted TB screening is warranted, especially among key populations, resulting in a greater yield of identifying TB cases, which aligns with the International Standards for Tuberculosis Care.

Tuberculosis symptom screening has limitations. However, there is a need to increase the frequency of when it is offered to MSM to enhance the opportunity of identifying those individuals in need of further investigations and treatment. Additionally, in Zimbabwe MSM experience considerable amounts of stigma and discrimination at most health facilities due to cultural imperatives as noted in the formative survey which was part of the BBS survey [7]. For Zimbabwe to ensure its health for all targets are met timeously there is a need to ensure that no group is left behind, including MSM and other key populations. A key intervention by the Zimbabwe Ministry of Health will be to train its healthcare workers in the provision of key population-friendly services and non-discriminatory behaviour.

Table 1 Baseline demographics and comparison of prevalence of TB symptoms among HIV Positive and HIV-negative MSM

Variable	HIV Positive	HIV negative	P-Value
Demographics			
Age in years			
Mean (Standard Deviation)	31.8 (9.2)	26.4 (7.9)	0.001
Ethnic Group			
Black African	327 (96.2)	1,151 (98.3)	0.035
Other races	13 (3.8)	20 (1.7)	
Education Level			
None	57 (4.9)	25 (7.4)	0.222
None and Primary	827 (70.6)	245 (72.1)	
Secondary	226 (19.3)	55 (16.2)	
Tertiary	61 (5.2)	15 (4.4)	
Vocational			
Employment Status			
Self-employed	103 (30.3)	262 (22.4)	0.001
Employed full time	59 (17.4)	149 (12.7)	
Employed part-time	35 (10.3)	123 (10.5)	
Full-time student	20 (5.9)	184 (15.7)	
Retired	123 (36.2)	453 (38.7)	
Unemployed and unemployed			
TB Symptoms			
In the last month, have you had a cough			
Prevalence	44/ 340 (40%)	62/1171 (5%)	0.001
In the last month, have you had night sweats?			
Prevalence	17/340 (5%)	37/1171 (3%)	0.108
In the last month, have you had any unplanned weight loss?			
Prevalence	43/340 (13%)	67/1171 (6%)	0.001
In the last month, have you had a fever or a "hot body"?			
Prevalence	9/340 (3%)	13/1171 (1%)	0.037
Experienced at least one of the four TB symptoms			
Prevalence	77/340 (23%)	125/1171 (11%)	0.001
In the past 12 months, have you been exposed to someone with TB?		. ,	
Prevalence	14/340 (4%)	39/1171 (3%)	0.487

Non-governmental organisations in Zimbabwe provide principally HIV-related services to MSM in secluded locals to avoid harassment from the police and security forces [7]. It will be also key to provide tuberculosis-related screening to those MSM using such services. Those individuals who screen positive for tuberculosis should immediately be provided with tuberculosis diagnostic tests and those testing positive should be immediately linked to care.

Abbreviations

- BBS Biobehavioral Survey
- HIV Human Immunodeficiency Virus
- MSM Men who have Sex with Men
- RDS Respondent Driven Sampling
- TB Tuberculosis

Acknowledgements

The authors would like to thank all individuals who took part in the survey.

Author contributions

GMus conceived the brief communication. GMus & MM carried out the statistical analysis. GMus & MM wrote the first draft of the brief communication. All authors (MM, IC, TD, BM, CS, DM, OM, EM, RB, GM, and GMus) contributed to the writing of the brief communication. All authors $(\mathsf{MM},\mathsf{IC},\mathsf{TD},\mathsf{BM},\mathsf{CS},\mathsf{DM},\mathsf{EM},\mathsf{RB},\mathsf{GM})$ read and approved the final brief communication.

Funding

The study was not funded.

Declarations

Competing interests

The authors declare no competing interests.

Received: 6 November 2023 / Accepted: 14 March 2024 Published online: 28 March 2024

References

- World Health Organization Global Tuberculosis Report. 2021. Geneva, Switzerland: WHO; 2021.
- World Health Organization Global Health Estimates. 2016: deaths by cause, age, sex, by country and by region. Geneva, Switzerland: WHO; 2018.
- Corbett EL, Marston B, Churchyard GJ, De Cock KM. Tuberculosis in sub-saharan Africa: opportunities, challenges and change in the era of antiretroviral treatment. Lancet. 2006;367:926–37.
- Borgdorff MW, Nagelkerke NJ, Dye C, Nunn P. Gender and tuberculosis: a comparison of prevalence surveys with notification data to explore sex differences in case detection. Int J Tuberculosis Lung Disease. 2000;4:123–32.

- Kemp JR, Gillian M, Simwaka BN, Salaniponi FM, Squire BS. Can Malawi's poor afford free tuberculosis services? Patient and household costs associated with a tuberculosis diagnosis in Lilongwe. Bull World Health Organ. 2007;85:580–5.
- Ministry of Health and Child Care (MoHCC). Zimbabwe Population-based HIV Impact Assessment (ZIMPHIA 2020): Final Report. Harare. 2021.
- HIV and STI Biobehavioral Survey among Men Who Have Sex with Men. Transgender women, and Genderqueer individuals in Zimbabwe - Final Report. New York: ICAP at Columbia University; 2020.
- Divala TH, Lewis J, Bulterys MA, Lutje V, Corbett EL, Schumacher SG, MacPherson P. Missed opportunities for diagnosis and treatment in patients with TB symptoms: a systematic review. Public Health Action. 2022;12(1):10–7. https://doi.org/10.5588/pha.21.0022.
- 9. UNAIDS. HIV prevention among key populations. 2016.
- Xin HN, Li XW, Zhang L, Li Z, Zhang HR, Yang Y. Tuberculosis infection testing in HIV-positive men who have sex with men from Xi'an China. Epidemiol Infect. 2017;145(3):498–502.
- 11. WHO, UNAIDS CDC, FHI 360. Biobehavioral survey guidelines for populations at risk for HIV. Geneva: World Health Organisation; 2017.
- 12. Storla DG, Yimer S, Bjune GA. A systematic review of delay in the diagnosis and treatment of tuberculosis. BMC Public Health. 2008;8(1):15.

- Kranzer K, Afnan-Holmes H, Tomlin K, Golub JE, Shapiro AE, Schaap A, Corbett EL, Lönnroth K, Glynn JR. The benefits to communities and individuals of screening for active tuberculosis disease: a systematic review. Int J Tuberc Lung Dis. 2013;17(4):432–46.
- Yoon C, Dowdy DW, Esmail H, MacPherson P, Schumacher SG. Screening for tuberculosis: time to move beyond symptoms. Lancet Respir Med. 2019;7(3):202–4.
- Willie B, Hakim AJ, Badman SG, Weikum D, Narokobi R, Coy K, Gabuzzi J, Pekon S, Gene S, Amos A, Kupul M, Hou P, Dala NM, Whiley DM, Wapling J, Kaldor KM, Vallely AJ, Kelly-Hanku A. High prevalence of pulmonary tuberculosis among female sex workers, men who have sex with men, and transgender women in Papua New Guinea. Trop Med Health. 2021;49:4. https://doi. org/10.1186/s41182-020-00293-w.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.